arXiv:1402.2303v2 [math.FA] 21 Apr 2014

# BANACH SPACES OF POLYNOMIALS AS "LARGE" SUBSPACES OF $\ell^\infty$-SPACES

ALEXANDER BRUDNYI

Abstract. In this note we study Banach spaces of traces of real polynomials on $\mathbb{R}^n$ to compact subsets equipped with supremum norms from the point of view of Geometric Functional Analysis.

## 1. Main Results

Recall that the *Banach-Mazur distance* between two $k$-dimensional real Banach spaces $E, F$ is defined as

$$d_{BM}(E,F) := \inf\{\|u\| \cdot \|u^{-1}\|\},$$

where the infimum is taken over all isomorphisms $u : E \to F$. We say that $E$ and $F$ are equivalent if they are isometrically isomorphic (i.e., $d_{BM}(E,F) = 1$). Then $\ln d_{BM}$ determines a metric on the set $\mathcal{B}_k$ of equivalence classes of isometrically isomorphic $k$-dimensional Banach spaces (called the *Banach-Mazur compactum*). It is known that $\mathcal{B}_k$ is compact of $d_{BM}$-"diameter" $\sim k$, see [G].

Let $C(K)$ be the Banach space of real continuous functions on a compact Hausdorff space $K$ equipped with the supremum norm. Let $F \subset C(K)$ be a filtered subalgebra with filtration $\{0\} \subset F_0 \subseteq F_1 \subseteq \cdots \subseteq F_d \subseteq \cdots \subseteq F$ (that is, $F = \cup_{d\in\mathbb{Z}_+} F_i$ and $F_i \cdot F_j \subset F_{i+j}$ for all $i, j \in \mathbb{Z}_+$) such that $n_d := \dim F_d < \infty$ for all $d$. In what follows we assume that $F_0$ contains constant functions on $K$. Our main result is

**Theorem 1.1.** *Suppose there are $c \in \mathbb{R}$ and $\{p_d\}_{d\in\mathbb{N}} \subset \mathbb{N}$ such that*

$$\frac{\ln n_{d\cdot p_d}}{p_d} \le c \quad \textit{for all} \quad d \in \mathbb{N}. \tag{1.1}$$

*Then there exist linear injective maps $i_d : F_d \hookrightarrow \ell^\infty_{n_{d\cdot p_d}}$ such that*

$$d_{BM}(F_d, i_d(F_d)) \le e^c, \quad d \in \mathbb{N}.$$

As a corollary we obtain:

**Corollary 1.2.** *Suppose $\{n_d\}_{d\in\mathbb{N}}$ grows at most polynomially in $d$, that is,*

$$\exists\, k, \hat{c} \in \mathbb{R}_+ \textit{ such that } \forall\, d \quad n_d \le \hat{c} d^k. \tag{1.2}$$

*Then for each natural number $s \ge 3$ there exist linear injective maps $i_{d,s} : F_d \hookrightarrow \ell^\infty_{N_{d,s}}$, where $N_{d,s} := \left\lfloor \hat{c} d^k \cdot s^k \cdot \left(\lfloor \ln(\hat{c} d^k)\rfloor + 1\right)^k \right\rfloor$, such that*

$$d_{BM}(F_d, i_{d,s}(F_d)) \le \left(e s^k\right)^{\frac{1}{s}}, \quad k \in \mathbb{N}.$$

2010 *Mathematics Subject Classification.* Primary 46B20, Secondary 46E15.
*Key words and phrases.* Banach-Mazur compactum, filtered algebra, entropy.
Research supported in part by NSERC.

Let $\mathcal{F}_{\hat{c},k}$ be the family of all possible filtered algebras $F$ on compact Hausdorff spaces $K$ satisfying condition (1.2). By $\mathcal{B}_{\hat{c},k,\bar{n}_d} \subset \mathcal{B}_{\bar{n}_d}$ we denote the closure in $\mathcal{B}_{\bar{n}_d}$ of the set formed by all subspaces $F_d$ of algebras $F \in \mathcal{F}_{\hat{c},k}$ having a fixed dimension $\bar{n}_d \in \mathbb{N}$.

Corollary 1.2 allows to estimate the metric entropy of $\mathcal{B}_{\hat{c},k,\bar{n}_d}$. Recall that for a compact subset $S \subset \mathcal{B}_{\bar{n}_d}$ its *$\varepsilon$-entropy* ($\varepsilon > 0$) is defined as $H(S,\varepsilon) := \ln N(S, d_{BM}, 1+\varepsilon)$, where $N(S, d_{BM}, 1+\varepsilon)$ is the smallest number of open $d_{BM}$-"balls" of radius $1+\varepsilon$ that cover $S$.

**Corollary 1.3.** *For $k \geq 1$ there exists a numerical constant $C$ such that for each $\varepsilon \in (0, \frac{1}{2}]$*

$$H\big(\mathcal{B}_{\hat{c},k,\bar{n}_d}, \varepsilon\big) \leq (Ck \cdot \ln(k+1))^k \cdot (\hat{c}d^k)^2 \cdot (\ln(\hat{c}d^k)+1)^{k+1} \cdot \left(\frac{1}{\varepsilon}\right)^k \cdot \left(\ln\left(\frac{1}{\varepsilon}\right)\right)^{k+1}.$$

## 2. Basic Example: Banach Spaces of Polynomials

Let $\mathcal{P}_d^n$ be the space of real polynomials on $\mathbb{R}^n$ of degree at most $d$. For a compact subset $K \subset \mathbb{R}^n$ by $\mathcal{P}_d^n|_K$ we denote the trace space of restrictions of polynomials in $\mathcal{P}_d^n$ to $K$ equipped with the supremum norm. Applying Corollary 1.2 to algebra $\mathcal{P}^n|_K := \cup_{d\geq 0} \mathcal{P}_d^n|_K$ we obtain:

(A) *There exist linear injective maps $i_{d,K} : \mathcal{P}_d^n|_K \hookrightarrow \ell^\infty_{N_{d,n}}$, where*

$$N_{d,n} := \big\lfloor e^{2n} \cdot (n+2)^{2n} \cdot d^n \cdot (2n+1+\lfloor n \ln d \rfloor)^n \big\rfloor, \tag{2.1}$$

*such that*

$$d_{BM}\left(\mathcal{P}_d^n|_K, i_{d,K}(\mathcal{P}_d^n|_K)\right) \leq \big(e \cdot (n+2)^2\big)^{\frac{1}{n+2}} \ (< 2.903). \tag{2.2}$$

Indeed,

$$\widetilde{N}_{d,n} := \dim \mathcal{P}_d^n|_K \leq \binom{d+n}{n} < \left(\frac{e\cdot(d+n)}{n}\right)^n \leq \left(\frac{e\cdot(1+n)}{n}\right)^n \cdot d^n < e^{2n} \cdot d^n. \tag{2.3}$$

Hence, Corollary 1.2 with $c = e^{2n}$, $k := n$ and $s := (n+2)^2$ implies the required result.

If $K$ is *$\mathcal{P}^n$-determining* (i.e., no nonzero polynomial vanish on $K$), then $\widetilde{N}_{d,n} = \binom{d+n}{n}$ and so for some constant $c(n)$ (depending on $n$ only) we have

$$\widetilde{N}_{d,n} < N_{d,n} \leq c(n) \cdot \widetilde{N}_{d,n} \cdot \big(1 + \ln \widetilde{N}_{d,n}\big)^n. \tag{2.4}$$

Hence, $V_{d,n}(K) := i_{d,K}(\mathcal{P}_d^n|_K)$ is a *"large" subspace* of $\ell^\infty_{N_{d,n}}$ in terminology of [B]. Therefore from (A) and [B, Prop. 8] applied to $V_{d,n}(K)$ we obtain:

(B) *There is a constant $c_1(n)$ (depending on $n$ only) such that for each $\mathcal{P}^n$-determining compact set $K \subset \mathbb{R}^n$ there exists an $m$-dimensional subspace $F \subset \mathcal{P}_d^n|_K$ with*

$$m := \dim F > c_1(n) \cdot \big(\widetilde{N}_{d,n}\big)^{\frac{1}{2}} \quad \text{and} \quad d_{BM}(F, \ell^\infty_m) \leq 3. \tag{2.5}$$

*In turn, if $\hat{d} \in \mathbb{N}$ is such that $N_{\hat{d},n} \leq c_1(n) \cdot \big(\widetilde{N}_{d,n}\big)^{\frac{1}{2}}$, then due to property (A) for each $\mathcal{P}^n$-determining compact set $K' \subset \mathbb{R}^n$ there exists a $\widetilde{N}_{\hat{d},n}$-dimensional subspace $F_{\hat{d},n,K'} \subset F$ such that*

$$d_{BM}\big(F_{\hat{d},n,K'}, \mathcal{P}_{\hat{d}}^n|_{K'}\big) < 9. \tag{2.6}$$

Further, the dual space $(V_d^n(K))^*$ of $V_d^n(K)$ is the quotient space of $\ell^1_{N_{d,n}}$. In particular, the closed ball of $(V_d^n(K))^*$ contains at most $c(n) \cdot \widetilde{N}_{d,n} \cdot \big(1 + \ln \widetilde{N}_{d,n}\big)^n$ extreme points, see (2.4). Thus the balls of $(V_d^n(K))^*$ and $V_d^n(K)$ are "quite different" as convex bodies. This is also expressed in the following property (similar to the celebrated John ellipsoid

theorem but with an extra logarithmic factor) which is a consequence of property (A) and [DMTJ, Prop. 1]:

(C) *There is a constant $c_2(n)$ (depending on $n$ only) such that for all $\mathcal{P}^n$-determining compact sets $K_1, K_2 \subset \mathbb{R}^n$*

$$d_{BM}\big(\mathcal{P}^n_d|_{K_1}, (\mathcal{P}^n_d|_{K_2})^*\big) \le c_2(n) \cdot (\widetilde{N}_{d,n} \cdot (1 + \ln \widetilde{N}_{d,n}))^{\frac{1}{2}}. \tag{2.7}$$

A stronger inequality is valid if we replace $(\mathcal{P}^n_d|_{K_2})^*$ above by $\ell^1_{\widetilde{N}_{d,n}}$, see [DMTJ, Th. 2].

**Remark 2.1.** Property (C) has the following geometric interpretation. By definition, $(\mathcal{P}^n_d|_{K_2})^*$ is a $\widetilde{N}_{d,n}$-dimensional real Banach space generated by evaluation functionals $\delta_x$ at points $x \in K_2$ with the closed unit ball being the balanced convex hull of the set $\{\delta_x\}_{x\in K_2}$. Thus $K_2$ admits a natural isometric embedding into the unit sphere of $(\mathcal{P}^n_d|_{K_2})^*$. Moreover, the Banach space of linear maps $(\mathcal{P}^n_d|_{K_2})^* \to \mathcal{P}^n_d|_{K_1}$ equipped with the operator norm is isometrically isomorphic to the Banach space of real polynomial maps $p : \mathbb{R}^n \to \mathcal{P}^n_d|_{K_1}$ of degree at most $d$ (i.e., $f^* \circ p \in \mathcal{P}^n_d$ for all $f^* \in (\mathcal{P}^n_d|_{K_1})^*$) with norm $\|p\| := \sup_{x\in K_2} \|p(x)\|_{\mathcal{P}^n_d|_{K_1}}$. Thus property (C) is equivalent to the following one:

(C′) *There exists a polynomial map $p : \mathbb{R}^n \to \mathcal{P}^n_d|_{K_1}$ of degree at most $d$ such that the balanced convex hull of $p(K_2)$ contains the closed unit ball of $\mathcal{P}^n_d|_{K_1}$ and is contained in the closed ball of radius $c_2(n) \cdot (\widetilde{N}_{d,n} \cdot (1 + \ln \widetilde{N}_{d,n}))^{\frac{1}{2}}$ of this space (both centered at 0).*

Our next property, a consequence of Corollary 1.3 and (2.3), estimates the metric entropy of the closure of the set $\widetilde{\mathcal{P}}_{d,n} \subset \mathcal{B}_{\widetilde{N}_{d,n}}$ formed by all $\widetilde{N}_{d,n}$-dimensional spaces $\mathcal{P}^n_d|_K$ with $\mathcal{P}^n$-determining compact subsets $K \subset \mathbb{R}^n$.

(D) *There exists a numerical constant $c > 0$ such that for each $\varepsilon \in (0, \frac{1}{2}]$,*

$$H\big(\mathrm{cl}(\widetilde{\mathcal{P}}_{d,n}), \varepsilon\big) \le (cn^2 \cdot \ln(n+1))^n \cdot d^{2n} \cdot (1 + \ln d)^{n+1} \cdot \left(\frac{1}{\varepsilon}\right)^n \cdot \left(\ln\left(\frac{1}{\varepsilon}\right)\right)^{n+1}. \tag{2.8}$$

**Remark 2.2.** The above estimate shows that $\widetilde{\mathcal{P}}_{d,n}$ with sufficiently large $d$ and $n$ is much less massive than $\mathcal{B}_{\widetilde{N}_{d,n}}$. Indeed, as follows from [Br, Th. 2]

$$H(\mathcal{B}_{\widetilde{N}_{d,n}}, \varepsilon) \sim \left(\frac{1}{\varepsilon}\right)^{\frac{\widetilde{N}_{d,n}-1}{2}} \quad \text{as} \quad \varepsilon \to 0^+$$

(here the equivalence depends on $d$ and $n$ as well). On the other hand, [P, Th. 1.1] implies that for any $\varepsilon > 0$,

$$0 < \liminf_{\widetilde{N}_{d,n}\to\infty} \frac{\ln H(\mathcal{B}_{\widetilde{N}_{d,n}}, \varepsilon)}{\widetilde{N}_{d,n}} \le \limsup_{\widetilde{N}_{d,n}\to\infty} \frac{\ln H(\mathcal{B}_{\widetilde{N}_{d,n}}, \varepsilon)}{\widetilde{N}_{d,n}} < \infty.$$

It might be of interest to *find sharp asymptotics of $H\big(\mathrm{cl}(\widetilde{\mathcal{P}}_{d,n}), \varepsilon\big)$ as $\varepsilon \to 0^+$ and $d \to \infty$*, and to *compute (up to a constant depending on $n$) $d_{BM}$-"diameter" of $\widetilde{\mathcal{P}}_{d,n}$*.

Similar results are valid for $K$ being a compact subset of a real algebraic variety $X \subset \mathbb{R}^n$ of dimension $m < n$ such that if a polynomial vanishes on $K$, then it vanishes on $X$ as well. In this case there are positive constants $c_X, \tilde{c}_X$ depending on $X$ only such that $\tilde{c}_X d^m \le \dim \mathcal{P}^n_d|_K \le c_X d^m$. For instance, Corollary 1.2 with $c = c_X$, $k := m$ and $s := (m+2)^2$ implies that $\mathcal{P}^n_d|_K$ is linearly embedded into $\ell^\infty_{N_{d,X}}$, where $N_{d,X} := \big\lfloor c_X d^m \cdot (m+2)^{2m} \cdot (\lfloor \ln(c_X d^m) \rfloor + 1)^m \big\rfloor$, with distortion $< 2.903$. We leave the details to the reader.

## 3. Proofs

*Proof of Theorem 1.1.* Since $\dim F_i = n_i$, $i \in \mathbb{N}$, and evaluations $\delta_z$ at points $z \in K$ determine bounded linear functionals on $F_i$, the Hahn-Banach theorem implies easily that $\operatorname{span}\{\delta_z\}_{z\in K} = F_i^*$. Moreover, $\|\delta_z\|_{F_i^*} = 1$ for all $z \in K$ and the closed unit ball of $F_i^*$ is the balanced convex hull of the set $\{\delta_z\}_{z\in K}$. Let $\{f_{1i}, \dots, f_{n_i i}\} \subset F_i$ be an Auerbach basis with the dual basis $\{\delta_{z_{1i}}, \dots, \delta_{z_{n_i i}}\} \subset F_i^*$, that is, $f_{ki}(\delta_{z_{li}}) := f_{ki}(z_{li}) = \delta_{kl}$ (the Kronecker-delta) and $\|f_{ki}\|_K = 1$ for all $k$. (Its construction is similar to that of the fundamental Lagrange interpolation polynomials for $F_i = \mathcal{P}_i^n|_K$, see, e.g., [BY, Prop. 2.2].)

Now, we use a "method of E. Landau" (see, e.g., [PS, Ch. 3, § 2]).

By the definition, for each $g \in F_i$ we have $g(z) = \sum_{k=1}^{n_i} f_{ki}(z)g(z_{ki})$, $z \in K$. Hence, $\|g\|_K \le n_i\|g\|_{\{z_{1i},\dots,z_{n_i i}\}}$. Applying the latter inequality to $g = f^{p_d}$, $f \in F_d$, containing in $F_i$, $i := d \cdot p_d$, and using condition (1.1) we get for $A_d := \{z_{1i}, \dots, z_{n_i i}\} \subset K$

$$\|f\|_K = (\|g\|_K)^{\frac{1}{p_d}} \le (n_{d\cdot p_d})^{\frac{1}{p_d}} \cdot (\|g\|_{A_d})^{\frac{1}{p_d}} \le e^c \cdot \|f\|_{A_d}.$$

Thus, restriction $F_d \mapsto F_d|_{A_d}$ determines the required map $i_d : F_d \hookrightarrow \ell^\infty_{n_{d\cdot p_d}}$. □

*Proof of Corollary 1.2.* We set $p_d := s \cdot (\lfloor \ln(\hat{c}d^k)\rfloor + 1)$, $d \in \mathbb{N}$. Then the condition of the corollary implies

$$\frac{\ln n_{d\cdot p_d}}{p_d} \le \frac{\ln(\hat{c}d^k) + k\ln p_d}{p_d} \le \frac{1}{s} + \frac{k\ln s}{s} =: c.$$

Thus the result follows from Theorem 1.1. □

*Proof of Corollary 1.3.* We make use of [P, Lm. 1.2] adapted to our setting:

**Lemma 3.1.** *Let $S_{\bar{n}_d} \subset \mathcal{B}_{\bar{n}_d}$ be the subset formed by all $\bar{n}_d$-dimensional subspaces of $\ell^\infty_{N_{d,s}}$. Consider $0 < \xi < \frac{1}{\bar{n}_d}$ and let $R = \frac{1+\xi\bar{n}_d}{1-\xi\bar{n}_d}$. Then $S_{\bar{n}_d}$ admits an $R$-net $T_R$ of cardinality at most $\left(1 + \frac{2}{\xi}\right)^{N_{d,s}\cdot\bar{n}_d}$.*

Now given $\varepsilon \in (0, \frac{1}{2}]$ we choose $s = \lfloor s_\varepsilon \rfloor$ with $s_\varepsilon$ satisfying $(es_\varepsilon^k)^{\frac{1}{s_\varepsilon}} = \sqrt[4]{1+\varepsilon}$ and $\xi$ such that $R = R_\varepsilon = \sqrt[4]{1+\varepsilon}$. Then according to Corollary 1.2 and Lemma 3.1, $\operatorname{dist}_{BM}\big(T_{R_\varepsilon}, \mathcal{B}_{\hat{c},k,\bar{n}_d}\big) < \sqrt{1+\varepsilon}$. For each $p \in T_{R_\varepsilon}$ we choose $q_p \in \mathcal{B}_{\hat{c},k,\bar{n}_d}$ such that $d_{BM}(p, q_p) < \sqrt{1+\varepsilon}$. Then the multiplicative triangle inequality for $d_{BM}$ implies that open $d_{BM}$-"balls" of radius $1+\varepsilon$ centered at points $q_p$, $p \in T_{R_\varepsilon}$, cover $\mathcal{B}_{\hat{c},k,\bar{n}_d}$. Hence,

$$N(\mathcal{B}_{\hat{c},k,\bar{n}_d}, d_{BM}, 1+\varepsilon) \le \operatorname{card} T_{R_\varepsilon} \le \left(1 + \frac{2}{\xi}\right)^{N_{d,s}\cdot\bar{n}_d}. \tag{3.9}$$

Next, the function $\varphi(x) = \ln(ex^k)^{\frac{1}{x}}$ decreases for $x \in [e^{\frac{k-1}{k}}, \infty)$ and $\lim_{x\to\infty}\varphi(x) = 0$. Its inverse $\varphi^{-1}$ on this interval has domain $(0, e^{-\frac{k-1}{k}}]$, increases and is easily seen (using that $\varphi \circ \varphi^{-1} = \mathrm{id}$) to satisfy

$$\varphi^{-1}(x) \le \frac{3k}{x} \cdot \ln\left(\frac{3k}{x}\right), \qquad x \in (0, e^{-\frac{k-1}{k}}].$$

Since $\frac{1}{4}\ln(1+\varepsilon) < e^{-\frac{k-1}{k}}$ for $\varepsilon \in (0, \frac{1}{2}]$, the required $s_\varepsilon$ exists and the previous inequality implies that

$$s_\varepsilon \le \frac{12k}{\ln(1+\varepsilon)} \cdot \ln\left(\frac{12k}{\ln(1+\varepsilon)}\right). \tag{3.10}$$

Further, we have

$$\frac{1}{\xi} = \frac{\bar{n}_d(1+R_\varepsilon)}{R_\varepsilon - 1} = \frac{\bar{n}_d(\sqrt[4]{1+\varepsilon}+1)}{\sqrt[4]{1+\varepsilon}-1} = \frac{\bar{n}_d(\sqrt[4]{1+\varepsilon}+1)^2\cdot(\sqrt{1+\varepsilon}+1)}{\varepsilon}. \tag{3.11}$$

From (3.9), (3.10), (3.11) invoking the definition of $N_{d,s}$ we obtain

$$\ln N(\mathcal{B}_{\hat{c},k,\bar{n}_d}, d_{BM}, 1+\varepsilon) \le \bar{n}_d\hat{c}d^k\left(\ln(\hat{c}d^k)+1\right)^k \ln\left(\frac{21\bar{n}_d}{\varepsilon}\right)\left(\frac{12k}{\ln(1+\varepsilon)}\ln\left(\frac{12k}{\ln(1+\varepsilon)}\right)\right)^k.$$

Using that $\bar{n}_d \le \hat{c}d^k$ and the inequality $\frac{2}{3}\cdot\varepsilon \le \ln(1+\varepsilon)$, $\varepsilon \in (0, \frac{1}{2}]$, we get the required estimate. □

## References


[B] J. Bourgain, Subspaces of $L_N^\infty$, arithmetical diameter and Sidon sets. Probability in Banach Spaces V. Proceed. Medford 1984. Lecture Notes in Math. **1153** (1985), 96–127.

[Br] E. M. Bronstein, $\varepsilon$-entropy of affine-equivalent convex bodies and Minkowski's compactum. (Russian) Optimizatsiya **39** No. 22 (1978), 5–11, 155.

[BY] A. Brudnyi and Y. Yomdin, Norming sets and related Remez-type inequalities. arXiv:1312.6050.

[G] E. D. Gluskin, The diameter of the Minkowski compactum is roughly equal to $n$. Funktsional. Anal. i Prilozhen. **15** (1) (1981), 72–73.

[DMTJ] W. J. Davis, V. D. Milman and N. Tomczak-Jaegermann, The distance between certain $n$-dimensional Banach spaces. Israel J. Math. **39** (1981), 1–15.

[P] G. Pisier, On the metric entropy of the Banach-Mazur compactum. arXiv:1306.5325v6.

[PS] G. Polya and G. Szegö, Problems and theorems in Analysis II. Springer, 1998.



Department of Mathematics and Statistics
University of Calgary
Calgary, Alberta
T2N 1N4
*E-mail address*: albru@math.ucalgary.ca